



UNIVERSITI PUTRA MALAYSIA

**DEVELOPMENT OF *IN VITRO* PROPAGATION TECHNIQUES FOR
ENDOSPERMUM MALACCENSE M.A AND *SHOREA PARVIFOLIA*
DYER**

AZIAH MOHD YUSOFF

FH 2003 13

**DEVELOPMENT OF *IN VITRO* PROPAGATION TECHNIQUES FOR
ENDOSPERMUM MALACCENSE M.A AND *SHOREA PARVIFOLIA* DYER**

By

AZIAH MOHD YUSOFF

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirements for the Degree of
Doctor of Philosophy**

April 2003



Abstract of thesis submitted to the Senate of University Putra Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy

**DEVELOPMENT OF *IN VITRO* PROPAGATION TECHNIQUES FOR
ENDOSPERMUM MALACCENSE M.A AND *SHOREA PARVIFOLIA* DYER**

by

AZIAH BINTI MOHD YUSOFF

April 2003

Chairman: Associate Prof. Dr. Nor Aini Ab. Shukor

Faculty: Forestry

Endospermum malaccense M.A and *Shorea parvifolia* Dyer are two commercially important timber species identified as potential plantation species. The procurement and storage of their seeds are difficult and is a major hindrance to plantation establishment. Development of micropropagation techniques is being pursued to provide an alternative in planting stock production.

Micropropagation of *E. malaccense* was achieved through *in vitro* production of plants through axillary shoots development from nodal segment explants of an elite tree. The explants were initially washed in 10 changes of sterile distilled water, followed by 5 changes of 0.05% (v/v) Tween 20 solution for 10 minutes each. This was then followed by rinsing in 10 changes of sterile distilled water and subsequently sterilised in a solution comprising 60 % (v/v) Clorox and 0.05% v/v Tween 20 for 10 minutes. After

which they were rinsed 10 times in 300ml sterile distilled water and finally immersed in 70% (v/v) ethanol for 1 minute. Shoots were induced on the nodal segment explants in MS basal salts supplemented with 22.2×10^{-6} M or 44.4×10^{-6} M. For shoot multiplication, the best medium is MS supplemented with 44.4×10^{-6} M BAP and solidified with a mixture of 1.7 g Gelrite and 4g Bacto agar per liter. *In vivo* rooting with Seradix 3 was more suitable for *in vitro* produced shoots of *E. malaccense*, compared with an *in vitro* rooting technique.

For *S. parvifolia*, high contamination was observed in all explant types. Multiple shoots were induced on nodal segments culture in WPM solid medium supplemented with 10^{-5} M BAP or 10^{-6} TDZ but were non-amenable to further subculture. Callus developed from immature seeds with gelatinous endosperm termed as embryonic masses in WPM supplemented with 10^{-4} M CPA induced callus formation. Globular shaped callus developed upon subculture. Histological examination of the globular shaped callus showed no evidence of somatic embryos formation. The globular structure was similar to the development of nodules.

The contaminants found on the immature seeds included a fungus, *Collectotrycum* spp. and a range of bacteria which are as follows: -*Kleibsellia planticola*, *Enterobacter agglomerans*, *Erwina* spp. (*E. uredora* or *E. herbicola*), *Serratia odorifera*, *Serratia marcescens*, *Serratia proteomaculans*, *Morganella morganii*, *Kluyera ascorbata*. A fern, *Asplenium nidus* was found contaminating the nodal segment cultures.

Abstrak tesis yang dikemukakan kepada Senat University Putra Malaysia sebagai memenuhi keperluan ijazah Doktor Falsafah

**PEMBENTUKAN TEKNIK MIKRO PERAMBATAN UNTUK
ENDOSPERMUM MALACCENSE M.A DAN *SHOREA PARVIFOLIA* DYER**

Oleh

AZIAH BINTI MOHD YUSOFF

April 2003

Pengerusi: Prof. Madya Dr. Nor Aini Ab. Shukor

Fakulti: Perhutanan

Endospermum malaccense M.A dan *Shorea parvifolia* Dyer, merupakan dua sepsis balak yang berpotensi ditanam secara ladang. Masalah pengutipan dan penyimpanan bijibenih menghindar kemajuan peladangan hutan secara komersial. Pembentukan teknik mikroperambatan dapat mengatasi masalah pembekalan bahan tanaman.

Mikroperambatan *E. Malaccense* tercapai apabila pembentukan pucuk aksil terbentuk ke atas eksplan bahagian nod pokok elit. Eksplan dibasmikuman dengan mencuci 10 kali dengan air suling steril, diikuti dengan 5 pencucian dalam larutan 0.05% (isipadu/isipadu) Tween 20 selama 10 minit, diikuti pula dengan 10 pertukaran air suling steril. Kemudian dicuci pula dalam larutan 60% (isipadu/isipadu) Clorox

dicampur dengan 0.05% (isipadu/isipadu) Tween 20 selama 10 menit. Kemudian dicuci dengan 10 kali dengan 300ml air suling steril dan akhir sekali direndam didalam 70% (isipadu/isipadu) etanol. Media MS dicampur dengan 22.2×10^{-6} M or 44.4×10^{-6} M sesuai dalam pengaruh pembentukan pucuk. Gandaan pucuk tercapai dalam campuran medium MS dan 44.4×10^{-6} M BAP yang dipejalkan dengan jel campuran 1.7 g Gelrite and 4g Bacto agar se liter. Teknik pengakaran *in vivo* dengan Seradix 3 lebih sesuai dibandingkan dengan teknik *in vitro*.

Untuk *S. parvifolia*, kadar kontaminasi tinggi dihadapi dalam kultur semua jenis eksplan. Pembentukan pucuk berbilang tercapai apabila bahagian nod di kulturkan didalam medium WPM yang ditambah dengan 10^{-5} M BAP atau 10^{-6} M TDZ. Pertambahan pucuk tidak tercapai dalam semua jenis medium. Pembentukan kalus terjadi ke atas eksplan endoperma cair di dalam medium WPM ditambah dengan 10^{-4} M CPA. Pengsubkultur menghasilkan kalus berbentuk globul. Kajian histology menunjukkan bahawa struktur ini adalah nodul.

Bijibenih muda dikontaminasikan oleh sejenis kulat, *Collectotrycum* spp. Dan pelbagai jenis bacteria seperti berikut:--*Kleibsellia planticola*, *Enterobacter agglomerans*, *Erwina* spp. (*E. uredora* or *E. herbicola*), *Serratia odorifera*, *Serratia marcesens*, *Serratia proteomaculans*, *Morganella morganii*, *Kluyera ascorbata*. Sejenis paku penumpang *Asplenium nidus* juga didapati menkontaminasikan kultur.

ACKNOWLEDGEMENTS

In the name of Allah S.W.T. The Most Gracious. The Most Merciful.

First and foremost I would like to convey my appreciation to my present employer the Golden Hope Research Sdn. Bhd. (GHR SB), for allowing me to continue my studies till completion. My appreciation goes to the Executive Director of Golden Hope Research Sdn. Bhd., Dr. Mohd Hashim Tajudin, for his support and to Dr. Mohd Nor Ghani, the Research Controller of the BioScience Division for his continuous encouragement. I am indebted to the management of the Forest Research Institute of Malaysia (FRIM), my former employer, for the permission to pursue my studies during my tenure in FRIM. Special thanks goes to the Director General Dato' Abd. Razak bin Mohd Ali, who has been very supportive of my pursuit to complete my studies.

I deeply appreciate the support of my supervisor Assoc. Prof. Dr. Nor Aini Shukor for her patience, encouragement and guidance. My heartfelt thanks go to the members supervisory committee, Assoc. Prof. Dr. Kamis Awang and Assoc. Prof. Dr. Midhzar Abdul Kadir for their suggestions and comments leading to the completion of this thesis.

To my former Research Assistant in FRIM, Pn. Sharifah Maulana, I thank her for assisting me in my experiments as well as her patience and fruitful suggestions. To my

present assistant at GHR SB, Pn. Ruziah Ahmad, I thank her for her assistance in the final layout and final touches of my thesis and to Asmida of FRIM, I am grateful for her assistance in the preparation of the colour prints for this thesis.

A million thanks goes to the members of the tissue culture laboratory at FRIM, Pn Fadhilah, Pn. Haliza, Pn. NorAsmah, Pn Halilah, Pn Normah, Pn Rozidah, Cik Sabariah, Pn Naemah, Pn Rukiah and Nizam for their cooperation.

To my former colleagues at FRIM, Dr. Ab. Rasip, Mohd Nor, David McKellar, Norhara, Dr. Ilham and numerous others, I thank you all for your support and suggestions.

To my beloved family, Mansor, my husband and children, Murni, Maizura, Marlina and Muhd. Isa, and last but not least to my mother, Cik An, my deepest appreciation, for their continuous encouragement, understanding and most of all, for their sacrifices in bearing all the shortcomings during these trying period, in order for me to complete this thesis. Thank you.

I certify that an Examination Committee met on 16th April 2003 to conduct the final examination of Aziah Mohd Yusoff on her Doctor of Philosophy thesis entitled “Development on *In Vitro* Propagation Techniques for *Endospermum Malaccense* M.A. and *Shorea Parvifolia* Dyer” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

JAMALUDDIN BASHARUDIN, Ph. D

Lecturer
Faculty of Forestry
University Putra Malaysia
(Chairman)

NOR AINI AB. SHUKOR, Ph. D

Associate Professor
Faculty of Forestry
University Putra Malaysia
(Member)

KAMIS AWANG, Ph. D

Associate Professor
Faculty of Forestry
University Putra Malaysia
(Member)

MIHDZAR ABDUL KADIR, Ph. D

Head
Department of Agriculture Technology
Faculty of Agriculture
University Putra Malaysia
(Member)

DATO A.H. ZAKRI, Ph. D

Professor
Institute of Advanced Studies
United Nations University
Tokyo, Japan
(Independent Examiner)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor / Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date : **16 JUN 2003**

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

NOR AINI AB. SHUKOR, Ph. D

Associate Professor
Faculty of Forestry
University Putra Malaysia
(Chairperson)

KAMIS AWANG, Ph. D

Associate Professor
Faculty of Forestry
University Putra Malaysia
(Member)

MIHDZAR ABDUL KADIR, Ph. D

Head
Department of Agriculture Technology
Faculty of Agriculture
University Putra Malaysia
(Member)



AINI IDERIS, Ph.D.

Professor / Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 11 JUL 2003

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



(AZIAH MOHD YUSOFF)

Date : 21ST MAY 2003

TABLE OF CONTENTS

		Page
	ABSTRACT.....	ii
	ABSTRAK.....	iv
	ACKNOWLEDGEMENTS	vi
	APPROVAL SHEETS.....	viii
	DECLARATION FORM.....	x
	TABLE OF CONTENTS.....	xi
	LISTS OF TABLES.....	xviii
	LISTS OF FIGURES.....	xxiii
	LISTS OF ABBREVIATIONS.....	xxix
	TERMINOLOGY.....	xxxix
 CHAPTER		
I	INTRODUCTION	1
 II	LITERATURE REVIEW	7
	<i>Shorea parvifolia</i> Dyer.....	7
	Botany and Taxonomy.....	7
	Distribution and Ecology.....	7
	Uses	7
	Propagation of <i>S. parvifolia</i>	8
	Potential as Plantation Species and the Improvement	8
	Activity	8
	<i>Endospermum malaccense</i> M.A.....	9
	Botany and Taxonomy.....	9
	Distribution and Ecology.....	9
	Uses	10
	Propagation of <i>E. malaccense</i>	10
	Status of Global Forest Plantation	10
	Plantation Forestry in Malaysia.....	12
	Plantation Establishment of Indigenous Species.....	14
	Planting Stock Production of Indigenous Species.....	16
	Vegetative Propagation.....	19
	<i>In Vitro</i> Propagation of Tropical Forest Species.....	21
	Explant Selection and Sterilisation Procedures.....	24
	Microbial Contaminants of Plant Tissue Culture.....	30
	Plant Production Through The Multiplication of Axillary	
	Shoots.....	32
	Factors Affecting Plant Production Through the Multiplication	
	of Axillary Shoots.....	33
	Genotype of Explants.....	33
	Media Composition.....	34
	Plant Growth Regulators.....	38
	Sugars as sources of carbohydrates.....	42
	Plant Production Through Somatic Embryogenesis.....	42
	Factors Affecting the Production and Development of	
	Embryogenic Callus.....	44

	Factors Associated with Explants.....	45
	Effects of Media Composition.....	48
	Effects of Plant Growth Regulators and Plant Growth Factors.....	50
	Effects of Carbohydrates Source.....	53
	Effects of Light.....	54
	Maturation and Germination of Somatic Embryos.....	55
	Effects of Absciscic Acid.....	56
	Effects of Carbohydrates.....	57
	Dessication.....	57
	Rooting and Establishment of Plantlets.....	58
	Media.....	59
	Sugars.....	59
	Plants Growth Regulators.....	60
	Light and Temperature.....	62
	Explant Support System.....	64
	Factors Associated with Explants.....	65
	Problems Faced in <i>In Vitro</i> Propagation of Plants.....	67
	Browning of Cultures.....	67
	Lag Time of Response from Explants of Mature Donors.....	68
	Vitrification of shoots	68
III	MATERIALS AND METHODS	70
	Introduction.....	70
	Preparation of Media	70
	Murashige and Skoog Media, MS (1962).....	70
	Gamborg B5 medium and Schenk and Hilderbrandt (SH) medium.....	70
	Woody Plant Medium, WPM (Lloyd and Mc Cown, 1980).....	71
	Preparation of Stock Solution of Growth Substances.....	71
	Description of Explants Used.....	72
	Identification of Contaminants.....	73
	Histological Techniques.....	75
	Resin Embedding.....	75
	Fixation.....	75
	Dehydration.....	77
	Infiltration.....	77
	Embedding.....	77
	Cutting.....	78
	Staining.....	79
	Mounting.....	80
	Preservation.....	81
	Microscopy.....	81

IV	MICROPROPAGATION OF <i>ENDOSPERMUM MALACCENSE</i>	82
	Introduction.....	82
	Establishment of Sterile Explants and Determination of Suitable Media Compositions.....	82
	Sterilisation of Explants.....	82
	Determination of suitable media for Establishment of Cultures	84
	Optimisation of Media for the Multiplication of Shoots	85
	Overcoming Vitrification.....	86
	Rooting Experiments and Acclimatisation of Shoots.....	87
	Results and Discussions.....	89
	Establishment of Sterile Explants and Determination of Suitable Media Compositions.....	89
	Sterilisation of Explants.....	89
	Determination of suitable media.....	90
	Multiplication of Shoots.....	91
	Overcoming Vitrification Problems	91
	Rooting And Acclimatisation Of Shoots.....	93
	Summary of Results.....	94
V	STUDIES TOWARDS <i>IN VITRO</i> PROPAGATION OF <i>SHOREA PARVIFOLIA</i>	95
	Introduction.....	95
	Explant Sources.....	95
	Mature Seeds.....	95
	<i>In Vitro</i> Germinated Seedlings.....	97
	Immature Seeds.....	98
	Explants from Seedlings Germinated in Non-Sterile Conditions.....	98
	Explants from Nursery Raised Seedlings.....	99
	Nodal segments and Shoot –Tips from Mature Trees.....	100
	Establishment of Sterile Culture and The Induction of Propagules On Various Explants of <i>S. parvifolia</i>	100
	Development of Sterilisation Protocol for Mature Seeds.....	100
	<i>In Vitro</i> Germination of Mature <i>S. parvifolia</i> Seeds.....	102
	Induction of Propagules on Explants Excised from <i>In Vitro</i> Germinated Seedlings.....	109
	Effects of MS and WPM Media in Combination with BAP and 2, 4-D on Explants Excised from <i>in vitro</i> Germinated Seedlings.....	109
	Effects of MS, WPM, SH and B5 Media in Combination with TDZ on Explants Excised from <i>in vitro</i> Germinated Seedlings.....	114

Establishment of Sterile Culture for Explants from Seedlings Germinated in Non-Sterile Conditions.....	125
Sterilisation Protocols and Media Used in Establishment of Sterile Culture for Explants from Seedlings in Non-sterile Conditions.....	125
Response of explants Excised from In Vivo Germinated Seedlings to Sterilisation Protocol and Media Used in the Induction of Propagagules.....	126
Establishment of Sterile Cultures for Shoot Tips and Nodal Segment Explants from Nursery Raised Seedlings.....	137
Development of Sterilisation Protocol for Shoot Tips and Nodal Segment Explants from Nursery Raised Seedlings.....	137
Effects of Different Sterilisation Protocols on Nodal Segments and Shoot Tip Explants from Nursery Raised Seedlings.....	140
Experiments with Fungicides and Antibiotic Pre- Treated Nursery Raised Seedlings.....	142
Pretreatment Protocols and Media Used for Explants from Pre-Treated Nursery Raised Seedlings.....	142
Response of Explants Excised from Nursery Raised Seedlings to Pre-Treatment.....	144
Response of Nodal Segments, Shoot Tip Explants Excised from Nursery-Raised Seedlings Pre-Treated with Fungicide and Antibiotic to Different Plant Growth Regulators.....	146
The Effect of Different Medium Formulations on Growth Responses of Nodal Segments and Shoot Tip Explants of <i>S. parvifolia</i> Cultured <i>in</i> <i>vitro</i>	149
The Effects of Supplementing BAP in WPM Medium on Nodal Segment and Leaf Explants Excised from Nursery Raised Seedlings.....	155
The Effects of Supplementing Different Combinations of BAP with Kinetin in WPM Medium on Nodal Segments and Leaf Explants Excised from Nursery Raised Seedlings.....	156
The Effects of Supplementing WPM Medium with TDZ on Shoot Tip and Nodal Segment Explants from Nursery Raised Seedlings.....	157
The Effects of Supplementing WPM Medium with NAA or 2,4-D on Explants Excised from Shoots of Nursery Raised Seedlings.....	159
Establishment of Sterile Cultures for Root Tip Explants from Nursery-Raised Seedlings.....	171

Sterilisation of Root Tip Explants from Nursery Raised Seedlings.....	171
Induction of Propagules in Root Tip Explants from Nursery Raised Seedlings.....	172
Establishment of Sterile Culture for Explants from Mature Trees.....	175
Sterilisation of Explants from Mature Trees.....	176
Induction of Propagules on Nodal Segments and Shoot Tips from Mature Trees.....	178
Establishment of Sterile Culture of Immature Seeds.....	
Sterilisation of Immature Seeds.....	180
Induction of Propagules on Explants Excised from Immature Seeds.....	184
Multiplication of Propagules.....	191
Multiplication of Propagules on Nodal Segment and Shoot Tip Explants Excised from <i>in vitro</i> and <i>in vivo</i> Germinated Seedlings.....	191
Multiplication of Shoots.....	191
Multiplication of Callus.....	197
Multiplication of Propagules on Nodal Segment and Shoot Tip Explant Excised from Nursery Raised Seedlings.....	200
Multiplication of Shoots.....	200
Multiplication of Callus.....	201
Callus Multiplication On Root Tip Explants.....	208
Effects of Several Different Types of PGRs on Callus Originating from Root Tips Explants.....	208
Effects of WPM Medium Supplemented with BAP and Kinetin on Callus Originating from Root Tips Explants.....	209
Effects of Sucrose, Abscissic Acid and Proline on Callus Originating from Root Tips Explants.....	216
Multiplication Of Callus On Immature Seeds Explants...	217
Contaminants Found in <i>S. parvifolia</i> Cultures.....	222
Fungal Contamination.....	222
Bacterial Contaminants.....	222
Contamination with Epiphytes.....	223
Histological Examination of Embryogenic Callus from Immature Seeds.....	227
Summary of Results.....	231

VI	DISCUSSIONS.....	237
----	------------------	-----

VII	CONCLUSIONS AND RECOMMENDATIONS	245
	REFERENCES.....	247
	APPENDICES.....	272
	VITA/BIODATA OF THE AUTHOR.....	304

LISTS OF TABLES

Table	Page
2.1 Annual plantation rates and plantation areas by region and species group	12
3.1 Tabulation of the plant growth regulators used in the experiments, their molecular weights, the desired quantity to prepare 50ml. Of 10^{-3} M stock solutions, their initial solvents and distilled water requirements	72
3.2 Glutaraldehyde-Paraformaldehyde-Caffeine fixative	75
3.3 Components of Phosphate buffer pH 7.2, 0.2 M	76
4.1 Effects of using four different sterilization protocol on nodal explants of <i>Endospermum maccalense</i> . Scoring was conducted as follows: +++=>61%, ++=31-60%, +=1-30%, - = 0%. Data were recorded 4 weeks after culture	90
4.2 Effects of the four different media formulations on development of nodal explants of <i>Endospermum maccalense</i> . Scoring was conducted as follows: +++ = >61%, ++=31-60%, +=1-30%, -+0%. Data were recorded 4 weeks after sub-culture	90
4.3 Summary of the t-Test on data from shoot multiplication of <i>Endospermum maccalense</i> in MS medium supplemented with either 2.22 or 4.44 x 10^{-6} M BAP	91
4.4 The effects of using different gelling agents on rates of vitrification of <i>Endospermum maccalense</i> shoots arising from nodal explants	93
4.5 The Effects of Commercial Rooting Powder Seradix 2 and Seradix 3 on in vitro Rooting and the Effects of the Solid MS Medium Supplemented with IBA and 10^{-6} M and 10^{-5} M on the <i>in vitro</i> Rooting of <i>E.malaccense</i> Shoots	93
5.1 Effects of different sterilization protocols on percentage browning, percentage germination, and percentage contamination of <i>Shorea parvifolia</i> seeds. Data were recorded after four weeks in culture. Values followed by the same letter within a column are not significantly different ($P<0.05$; Duncan's Multiple Range Test)	102

5.2	Effects of different media on percentage browning, percentage germination, and percentage contamination of <i>Shorea parvifolia</i> seeds. Observations were made after four week in culture. Values followed by the same letter within a column are not significantly different ($P<0.05$; Duncan's Multiple Range Test)	108
5.3	Number of hypocotyls, leaf, nodal segment, root and shoot explants cultured in either MS or WPM basal medium, regardless of the combination of BAP and 2,4-D used	110
5.4	Responses of explants from <i>in vitro</i> seedlings of <i>Shorea parvifolia</i> cultured on Murashige and Skoog medium containing benzylamino purine (at 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M) alone, or in combination with 2,4-dichlorophenoxyacetic acid (at 10^{-4} M) after 4 weeks in culture	117
5.5	Responses of explants from <i>in vitro</i> germinated seedlings of <i>Shorea parvifolia</i> cultured on Woody Plant Medium (1980) containing benzylamino purine (at 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M) alone, or in combination with 2,4-dichlorophenoxyacetic acid (at 10^{-4} M) after 4 weeks in culture	119
5.6	Responses of explants from <i>in vitro</i> germinated seedlings of <i>Shorea parvifolia</i> cultured on Murashige and Skoog MS (1962). Schenk and Hiderbrandt, SH (1972), Gamborg et al, B5, (1968) and Woody Plant Medium (WPM), (Llyod and Mc Cown, 1980) in combination with 10^{-8} M to 10^{-6} M thidiazuron (TDZ) after 4 weeks in culture	122
5.7	Responses of explants from <i>in vivo</i> germinated seedlings of <i>Shorea parvifolia</i> cultured on Murashige and Skoog medium containing benzylamino purine (at 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M) alone, or in combination with 2,4-dichlorophenoxyacetic acid (at 10^{-4} M) after 4 weeks in culture	129

5.8	Responses of explants from <i>in vivo</i> germinated seedlings of <i>Shorea parvifolia</i> cultured on Woody Plant Medium containing benzylamino purine (at 10^{-4} M, 10^{-5} M, 10^{-6} M, 10^{-7} M) alone, or in combination with 2,4-dichlorophenoxyacetic acid (at 10^{-4} M) after 4 weeks in culture	133
5.9	Effects of different sterilization protocols on nodal and shoot tip explants excised from nursery raised seedlings of <i>Shorea parvifolia</i> . Percentage contamination was observed after 4 weeks in WPM medium	141
5.10	Effects of pre-treatment of <i>Shorea parvifolia</i> nursery raised seedlings by root drenching and/or spraying with antibiotics (streptomycin and tetracycline) alone or in combination with the fungicide Benlate. Seedlings were treated once a week for 1, 2, 3, or 4 weeks, before excision of nodal segments, leaves and shoot tips. Observations were made after 4 weeks in culture on WPM medium supplemented with TDZ, BAP, kinetin, NAA or 2,4-d alone	146
5.11	Effects of different concentrations of 2,4-D, NAA, BAP, Kinetin and thidiazuron (TDZ) on callus formation, shoot elongation and browning of leaf, nodal segments and shoot tip explants excised from nursery raised seedlings of <i>Shorea parvifolia</i>	149
5.12	Response of explants of <i>Shorea parvifolia</i> explants from nursery raised seedlings to different basal medium compositions. Data were recorded after 4 weeks in culture. (There was no significance difference at $\alpha=0.05$ for all the means tested)	151
5.13	Response of explants of <i>Shorea parvifolia</i> explants from nursery raised seedlings to WPM medium supplemented with 10^{-4} M, 10^{-5} M, 10^{-6} M and 10^{-7} M BAP. Data were recorded after 4 weeks in culture. (There was no significance difference at $\alpha=0.05$ for all the means tested)	156

5.14	Responses of explants of <i>Shorea parvifolia</i> explants from nursery raised seedlings to WPM medium supplemented with BAP(10^{-7} M to 10^{-4} M) in combination with kinetin (10^{-5} M to 10^{-7} M). Data were recorded after 4 weeks in culture. (There was no significance difference at $\alpha=0.05$ for all the means tested)	157
5.15	Response of <i>Shorea parvifolia</i> nodal segments and shoot tip explants excised from nursery raised seedlings to WPM medium supplemented with TDZ (10^{-9} M to 10^{-5} M). Data were recorded after 4 weeks in culture. (There was no significance difference at $\alpha=0.05$ for all the means tested)	158
5.16	Response of all explant types (mid-rib, nodal segment, leaf, petiole and shoot tips) from nursery raised seedlings to WPM medium supplemented with 10^{-7} M to 10^{-4} M 2,4-D. (There was no significance at $\alpha=0.05$ for all the means tested)	160
5.17	Response of leaf, mid-rib, nodal segment, petiole and shoot tip explants initially cultured in WPM media supplemented with 2,4-D at concentrations (10^{-7} M to 10^{-4} M) to WPM medium supplemented with 10^{-7} M to 10^{-8} M 2,4-D	164
5.18	Response of all explant types (mid-rib), nodal segment, leaf, petiole and shoot tips) from nursery raised seedlings to WPM supplemented with 10^{-7} M to 10^{-4} M NAA	168
5.19	Response of leaf, mid-rib, nodal segment, petiole and shoot tip explants initially cultured in WPM media supplemented with NAA at concentrations (10^{-7} M to 10^{-4} M) to WPM medium supplemented with 10^{-7} M to 10^{-8} M NAA	170
5.20	Results of the effect different concentration of Mercuric chloride solution on root explants in terms of non-contamination and response. The different concentration of mercuric chloride used were S1=0.1% (w/v), S2=0.2% (w/v), S3=0.3% (w/v). (There was no significance difference at $\alpha=0.05$ for all the means tested)	172

5.21	Results of the effect of supplementing 2,4-D or NAA to Woody Plant Medium on root explants excised from nursery grown seedling. (There was no significance difference at $\alpha=0.05$ for all the means tested)	175
5.22	Results of different treatments used during collection of <i>Shorea parvifolia</i> explants from a 40-year old tree within FRIM campus. Shoot tips and nodal segment were cultured in half strength MS, half-strength WPM, WPM (NH ₄ NO ₃), SH and B5 media. Explants were sterilized according to method 1 (page 173)	178
5.23	Responses of nodal segments and shoot tip explants of <i>Shorea parvifolia</i> to three sterilization protocols when cultured on half-strength Murashige and Skoog medium (MS), half-strength Woody Plant Medium (WPM), WPM lacking ammonium nitrate, Shenck and Hildebrandt Medium (SH) or Gamborg's B5 medium (B5) after 4 weeks in culture	180
5.24	Response of immature <i>S. parvifolia</i> seeds to four different sterilization treatments and cultured in media listed on page 72, four weeks after culture. (There was no significance difference at $\alpha=0.05$ for all the means tested)	183
5.25	Types of Plant Growth Regulators Used and Their Abbreviations	184
5.26	Response of cotyledon, embryonic mass and whole seeds* of <i>S. parvifolia</i> to WPM medium supplemented with 10 ⁻⁷ to 10 ⁻⁴ to several different types of Plant Growth Regulators (PGR) added alone after 4 weeks in culture	187
5.27	Results on multiplication of shoots induced on nodal segments and shoot tip explants excised from <i>in vitro</i> and <i>in vivo</i> germinated seedlings of <i>S. parvifolia</i> cultured in WPM medium supplemented with BAP (10 ⁻⁵ M –10 ⁻⁷ M) or TDZ (10 ⁻⁹ M-10 ⁻⁵ M). Observation was carried out four weeks after culture. For each treatment six explants were used	192

5.28	Results on multiplication of callus induced on leaf, cotyledon, hypocotyle, nodal segment and shoot tip explants excised from <i>in vitro</i> and <i>in vivo</i> germinated seedlings of <i>S. parvifolia</i> cultured in WPM medium supplemented with 2,4-D (10^{-4} M – 10^{-7} M) or TDZ (10^{-8} M- 10^{-4} M). Observation was carried out six weeks after culture	198
5.29	The number of tubes used per treatment in callus multiplication experiment. The callus were classified according to the explant origin of callus	201
5.30	Effects of solid WPM medium supplemented with 2,4-D, NAA, BAP, kinetin, NOA, 4-CPA, MCPA, Dicamba and Picloran at levels (10^{-6} M to 10^{-4} M) after six weeks in culture	210
5.31	Effects of solid WPM medium supplemented with a combination of BAP and Kinetin on callus originating from roots of <i>S. parvifolia</i> six weeks after culture. Both PGRs were used at 10^{-7} M, 10^{-6} M and 10^{-5} M	213
5.32	Effects of different levels of Absciscic acid (ABA), sucrose and proline on callus originating from roots	217
5.33	Effects of 2,4-D, NAA, Dicamba, MCPA, NOA, 4-CPA and Picloram on callus originating from embryos	219

LIST OF FIGURES

Figure	Page
2.1 Different phases in somatic embryogenesis (Bornman, 1993)	45
4.1 <i>In vitro</i> rooting of <i>Endospermum malaccense</i> by dipping shoots bases in either Seradix 2 or Seradix 3 and placing shoots in seed trays containing sand, with 10 shoots per tray	88
4.2 <i>In vitro</i> Rooting of <i>Endospermum malaccense</i> in half-strength MS medium solidified with mixture of Bacto-agar and Gelrite and supplemented with either 10^{-5} M IBA or 10^{-6} M IBA	88
5.1 Mature seeds of <i>Shorea parvifolia</i>	96
5.2 <i>In vitro</i> germination of <i>S. parvifolia</i> seeds in liquid medium with filter paper support which was prepared by folding two pieces 15 cm disc of Whatman No.1 filter paper folded length-wise. Each fold was 1 cm wide and wavelike fan folds was created and the seeds were placed in between the folds	104
5.3 Response of Hypocotyl, Leaf, Nodal segments and Root Explants Originating from <i>in vitro</i> germinated seedlings after 4 weeks in WPM and MS media	112
5.4 Response of Hypocotyl, Leaf, Nodal segments Explants Originating from <i>in vivo</i> germinated seedlings after 4 weeks in MS and WPM media	128
5.5 Formation of callus on leaf explants excised from <i>S. parvifolia</i> nursery raised seedlings four weeks after culture in 10^{-7} M TDZ	148

5.6	Response of shoot tip explants excised from nursery raised seedlings of <i>S. parvifolia</i> to different media basal used after 4 weeks in culture. (B5 = Gamborg Medium (1968), MS = Murashige & Skoog (1962), SH = Schenk Hilderbrandt (1972), WPM = Woody Plant Medium (Lloyd & Mc Cown, 1980), wpm- = WPM = Woody Plant Medium (Lloyd & Mc Cown, 1980) without NH_4NO_3)	153
5.7	Response of nodal segment explants excised from nursery raised seedlings of <i>S. parvifolia</i> to different basal media used after 4 weeks in culture. (B5 = Gamborg Medium (1968), MS = Murashige & Skoog (1962), SH = Schenk Hilderbrandt (1972), WPM = Woody Plant Medium (Lloyd & Mc Cown, 1980), wpm- = WPM = Woody Plant Medium (Lloyd & Mc Cown, 1980) without NH_4NO_3)	154
5.8	Response of leaf, petiole, nodal segment (ns), shoot-tip (st) and mid-rib explants to WPN medium supplemented with 10^{-7} M to 10^{-4} M 2, 4-D after 4 weeks in culture. ($\text{D}_4 = 10^{-4}$ M 2, 4-D, $\text{D}_5 = 10^{-5}$ M 2, 4-D, $\text{D}^6 = 10^{-6}$ M 2, 4-D and $\text{D}_7 = 10^{-7}$ M 2, 4-D)	163
5.9	Roots developed on callus formed on nodal segment explants. Callus was induced on nodal segment explants in media supplemented 10^{-4} M 2, 4-D after six weeks in culture. Development of roots were observed after transfer to media supplemented with 10^{-8} M 2, 4-D	165
5.10	Development of roots on nodal segment explants excised from nursery raised germinated seedlings of <i>S. parvifolia</i> , when the explant were transferred WPM medium supplemented with 10^{-4} M 2, 4-D to WPM medium containing 10^{-7} M 2, 4-D	166
5.11	Development of roots on nodal segment explants excised from nursery raised germinated seedlings of <i>S. parvifolia</i> , when the explant were transferred WPM medium supplemented with 10^{-4} M 2, 4-D to WPM medium containing 10^{-7} M 2, 4-D	166